REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of Information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed; and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA :22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (10704-0188), Washington, DC 20503.

AGENCY USE ONLY (Leave blank) 2. REPORT DATE 3. REPORT TYPE AND	D DATES COVERED
12-20-05 Final	12/01/00 - 9/30/05
TITLE AND SUBTITLE	S. FUNDING NUMBERS
COAMPS Simulations of the Coastal Atmosphere	l
AUTHOR(S) Roger M. Samelson	N00014-01-1-0231
Eric D. Skyllingstad	
PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION
Corvalls, OR 97331	REPORT NUMBER
Corvalls, OR 97331	NO1450
. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)	10. SPONSORING/MONITORING
ONR	AGENCY REPORT NUMBER
1. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION/AVAILABILITY STATEMENT	126. DISTRIBUTION CODE
Unlimited Public Access DISTRIBUTION STATEMENTA	
Approved for Public Release	
13. ABSTRACT (Maximum 200 words) Work has focused primarily on simulations of the coastal atmospher coupled regional ocean-atmosphere model, based on the Naval Rese COAMPS and Rutgers/UCLA ROMS models, for use on multiple processors. Progress to date includes the development of a processor of module using the Model Coupling Toolkit (MCT) (Argonne National management module is able to run both COAMPS and ROMS on in processors. Model initializations for both the ocean and atmosphere for an idealized, linear coastline, using conditions similar to summer along the U.S. west coast.	arch Laboratory rocessor, cluster nanagement code al Laboratory). The dependent
Comparison of the coupled simulations with uncoupled and partial suggests that the dynamics of the near shore upwelling region are coupled processes. For example, in cases without coupling, offsh nearly constant across the coastal shelf. Coupling generates a transport results that gradually increases over a distance of 10-20 shore currents are also affected with the coastal jet decreasing near wind stress is reduced. 14. SUBJECT TERMS	re greatly affected by nore mass transport is a more realistic mass
	16. PRICE CODE
17. SECURITY CLASSIFICATION 18. SECURITY CLASSIFICATION 19. SECURITY CLASSIFICATION OF ABSTRACT	SIFICATION 20. LIMITATION OF ABSTR

FINAL TECHNICAL REPORT

ONR GRANT #N00014-01-1-0231

PIs: Eric D. Skyllingstad Roger M. Samelson

Title: COAMPS Simulations of the Coastal Atmosphere

Work has focused primarily on simulations of the coastal atmosphere, and on building a coupled regional ocean-atmosphere model, based on the Naval Research Laboratory COAMPS and Rutgers/UCLA ROMS models, for use on multiple processor, cluster systems. Progress to date includes the development of a processor management code module using the Model Coupling Toolkit (MCT) (Argonne National Laboratory). The management module is able to run both COAMPS and ROMS on independent processors. Model initializations for both the ocean and atmosphere have been developed for an idealized, linear coastline, using conditions similar to summer upwelling events along the U.S. west coast.

Results from our experiments focus mostly on idealized upwelling case for examining the coupled response of the upwelling system and providing a test for the coupled COAMPS/ROMS model that is under development. For these studies, we have chosen a simple channel domain with a linear coast line, no topography on land, a constant ocean shelf slope, and periodic conditions in the north-south direction. Our coupled simulations examine how changes in the SST forced by upwelling wind conditions affect the lower atmospheric boundary layer and surface stress. We find that cooling of the atmosphere by upwelled water causes a decoupled boundary layer structure. The net effect is a reduction in the wind stress by about a factor of 2 within 10 km of the coast, with an accompanying increase in the wind stress curl.

Comparison of the coupled simulations with uncoupled and partial coupled cases suggests that the dynamics of the near shore upwelling region are greatly affected by coupled processes. For example, in cases without coupling, offshore mass transport is nearly constant across the coastal shelf. Coupling generates a more realistic mass transport results that gradually increases over a distance of 10-20 km off shore. Near shore currents are also affected with the coastal jet decreasing near the coast where the wind stress is reduced.

DISTRIBUTION STATEMENT A

Approved for Public Release
Distribution Unlimited

20051227 040

LIST OF PUBLICATIONS

- Perlin, N., R. M. Samelson, and D. B. Chelton, 2004. Scatterometer and model wind and wind stress in the Oregon-California coastal zone. Monthly Weather Review, 132, 2110-2129.
- Samelson, R. M., E. D. Skyllingstad, D. B. Chelton, S. K. Esbensen, L. W. O'Neill, and N. Thum, 2005. A note on the coupling of wind stress and sea surface temperature. Journal of Climate, accepted.
- Skyllingstad, E. D., R. Samelson, L. Mahrt, P. Barbour, 2004: A numerical modeling study of warm offshore flow over cool water. Monthly Weather Review, 133, 345-361.
- Perlin, N., E. Skyllingstad, R. Samelson, and P. Barbour, 2006: A numerical study of idealized upwelling conditions using a coupled mesoscale ocean-atmosphere model. In preparation.